



The slow ape: High infant survival and long interbirth intervals in wild orangutans

van Noordwijk, Maria A ; Utami Atmoko, S Suci ; Knott, Cheryl D ; Kuze, Noko ; Morrogh-Bernard, Helen C ; Oram, Felicity ; Schuppli, Caroline ; van Schaik, Carel P ; Willems, Erik P

Abstract: Orangutans (*Pongo* spp.) are reported to have extremely slow life histories, including the longest average interbirth intervals of all mammals. Such slow life history can be viable only when unavoidable mortality is kept low. Thus, orangutans' survivorship under natural conditions is expected to be extremely high. Previous estimates of orangutan life history were based on captive individuals living under very different circumstances or on small samples from wild populations. Here, we combine birth data from seven field sites, each with demographic data collection for at least 10 years (range 12–43 years) on wild orangutans to better document their life history. Using strict criteria for data inclusion, we calculated infant survival, interbirth intervals and female age at first reproduction, across species, subspecies and islands. We found an average closed interbirth interval of 7.6 years, as well as consistently very high pre-weaning survival for males and females. Female survival of 94% until age at first birth (at around age 15 years) was higher than reported for any other mammal species under natural conditions. Similarly, annual survival among parous females is very high, but longevity remains to be estimated. Current data suggest no major life history differences between Sumatran and Bornean orangutans. The high offspring survival is remarkable, noting that modern human populations seem to have reached the same level of survival only in the 20th century. The orangutans' slow life history illustrates what can be achieved if a hominoid bauplan is exposed to low unavoidable mortality. Their high survival is likely due to their arboreal and non-gregarious lifestyle, and has allowed them to maintain viable populations, despite living in low-productivity habitats. However, their slow life history also implies that orangutans are highly vulnerable to a catastrophic population crash in the face of drastic habitat change.

DOI: <https://doi.org/10.1016/j.jhevol.2018.09.004>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-166291>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

van Noordwijk, Maria A; Utami Atmoko, S Suci; Knott, Cheryl D; Kuze, Noko; Morrogh-Bernard, Helen C; Oram, Felicity; Schuppli, Caroline; van Schaik, Carel P; Willems, Erik P (2018). The slow ape: High

infant survival and long interbirth intervals in wild orangutans. *Journal of Human Evolution*, 125:38-49.
DOI: <https://doi.org/10.1016/j.jhevol.2018.09.004>

Accepted for Publication 3 October 2018

Journal of Human Evolution 125 (2018) 38-49

<https://doi.org/10.1016/j.jhevol.2018.09.004>

The slow ape: high infant survival and long interbirth intervals in wild orangutans

Maria A. van Noordwijk¹, S. Suci Utami Atmoko², Cheryl D. Knott³, Noko Kuze⁴, Helen C.

Morrogh-Bernard⁵, Felicity Oram⁶, Caroline Schuppli¹, Carel P. van Schaik¹, Erik P. Willems¹

Affiliations:

¹ Department of Anthropology, University of Zürich, Winterthurerstr 190, 8057 Zürich, Switzerland

² Fakultas Biologi, Universitas Nasional, Jln Sawo Manila, Jakarta 12520, Indonesia

³ Department of Anthropology, Boston University, Boston, MA 02215 USA

⁴ Department of Anthropology, The National Museum of Nature and Science, Ibaraki 305-0005, Japan

⁵ Borneo Nature Foundation, www.borneonature.org, and College of Life and Environmental Science, University of Exeter, Cornwall, TR10 9FE, England, UK

⁶ Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah, Kota Kinabalu, Sabah 88400, Malaysia and HUTAN–Kinabatangan Orangutan Conservation Programme Sandakan, Sabah, 88999 Malaysia

Corresponding author: Maria A. van Noordwijk

Email: vnoord@aim.uzh.ch

Phone +41 44 635 5436

Declarations of interest: none

ABSTRACT

Orangutans (*Pongo* spp.) are reported to have extremely slow life histories, including the longest average interbirth intervals of all mammals. Such slow life history can be viable only when unavoidable mortality is kept low. Thus, orangutans' survivorship under natural conditions is expected to be extremely high. Previous estimates of orangutan life history were based on captive individuals living under very different circumstances or on small samples from wild populations. Here, we combine birth data from seven field sites, each with demographic data collection for at least 10 years (range 12-43 years) on wild orangutans to better document their life history. Using strict criteria for data inclusion, we calculated infant survival, interbirth intervals and female age at first reproduction, across species, subspecies and islands. We found an average closed interbirth interval of 7.6 years, as well as consistently very high pre-weaning survival for males and females. Female survival of 94% until age at first birth (at around age 15 years) was higher than reported for any other mammal species under natural conditions. Similarly, annual survival among parous females is very high, but longevity remains to be estimated. Current data suggest no major life history differences between Sumatran and Bornean orangutans. The high offspring survival is remarkable, noting that modern human populations seem to have reached the same level of survival only in the 20th century. The orangutans' slow life history illustrates what can be achieved if a hominoid bauplan is exposed to low unavoidable mortality. Their high survival is likely due to their arboreal and non-gregarious lifestyle, and has allowed them to maintain viable populations, despite living in low-productivity habitats. However, their slow life history also implies that orangutans are highly vulnerable to a catastrophic population crash in the face of drastic habitat change.

Keywords: life history; first birth; interbirth interval; immature survival; great ape

INTRODUCTION

Primates, and in particular Hominoidea, are characterized by a slow life history compared to other mammals (Stearns, 2000; van Schaik and Isler, 2012). They usually produce single offspring that mature slowly, have a late age at first reproduction (AFR) and a relatively long lifespan. All hominids basically show slow life histories regardless of living conditions (Knott 2001). Indeed, some recent studies that compared the life histories of wild chimpanzees and nomadic human foragers (e.g., Muller and Wrangham, 2014; Blurton Jones, 2016; Wood et al., 2017) concluded that the differences were much smaller than previously estimated (Hill et al., 2001).

Such slow development and low reproductive rates are viable only when the extrinsic risk of mortality is low throughout the lifespan (Stearns 2000). In general, pre-senescence, ‘unavoidable’ mortality is mainly attributed to predation, starvation or disease. However, at present, we can only tentatively link varying unavoidable mortality to variation in life histories among the extant great apes (Knott 2001). The observed faster life history of gorillas compared to chimpanzees (Knott, 2001; Bronowski et al., 2016a, b) is correlated with folivory, although why (and even whether) folivory generally speeds up life-history pace is still unclear (Janson and van Schaik, 1993; Borries et al., 2013). One would expect orangutans and chimpanzees to be most similar among the apes, given their similar female body size as well as ecological similarity as extractively foraging frugivores, even though they occupy different geographic regions (Asia versus Africa). However, available evidence suggested that orangutans have slower life histories

than chimpanzees and indeed all other primates (Knott, 2001; Wich et al., 2004, 2009a; Marshall et al., 2009b). In particular, orangutans appeared to have extremely long IBIs (interbirth intervals; e.g., Galdikas and Wood, 1990; Knott, 2001; Wich et al., 2004). Even when provisioned with a steady, predictable food supply and veterinary care, captive orangutan IBIs, at around 5.5 years (Anderson et al., 2008; Shumaker et al., 2008), tend to be longer than among other apes (Knott, 2001). However, sample sizes for orangutan life-history parameters under natural conditions have remained small (Wich et al., 2004), and here we revisit this preliminary conclusion based on a more extensive dataset.

Among the suggested reasons for this especially slow life history, low forest productivity may be most relevant (Knott, 2001, 2004). Compared to African forests, the fruit production in Southeast Asian forests is lower (Terborgh and van Schaik 1987). Lower productivity may translate into lower foraging yields, which in turn may constrain reproduction and growth (Pontzer and Kamilar, 2009). Forest productivity is also less predictable in Southeast Asia, with widely spaced periods of habitat-wide high peaks, so-called ‘mast-fruiting’, and multi-year periods of variable duration with very low fruit availability (Ashton et al., 1988; van Schaik and Pfannes, 2005; Hanya et al., 2011). The latter may lead to periods of severe scarcity (Knott, 1998), potentially resulting in increased unavoidable mortality. On the other hand, it has been argued that both very slow growth (Janson and van Schaik 1993) and an extended period of low maternal nutritional support buffer against starvation of both mothers and ecologically less competent immatures (Janson and van Schaik, 1993; van Noordwijk et al., 2013).

The lower food availability in Asia than Africa has also been suggested to impose a less gregarious life style on the Asian great apes (van Schaik, 1999; Wich et al., 1999). In addition, orangutans are much more arboreal than the African apes. Arboreality itself may reduce

predation risk, a major source of unavoidable mortality, and therefore allow for an even slower life history, within both primates and mammals in general (van Schaik and Deaner, 2003). The combination of non-gregariousness and arboreality, in turn, may largely protect orangutans against transmission of contagious diseases, responsible for some immature deaths in chimpanzees (Kuehl et al., 2008; Hassel et al., 2017) and orphaned orangutans in rehabilitation centers (Galdikas and Ashbury, 2012; Kuze et al., 2012). It also limits exposure to soil-dwelling parasites, especially detrimental to younger animals, and transmission of primate and human diseases that were implicated as causal factors in the lower survival of orangutans at rescue centers (Mul et al., 2007, Foitova et al., 2009). Finally, arboreality may impose an additional burden on mothers who carry their offspring in the trees and need to constantly protect them against falling (e.g., Pontzer and Wrangham, 2004; Halsey et al., 2016), thus possibly further slowing down a mother's reproductive rate (Knott, 2001, 2004). For example, in human hunter-gatherers the cost of infant carrying is suggested to delay a mother's investment in a subsequent child (Blurton-Jones 1986; Kramer 1998).

To date, the net effect of all these potential factors on the Asian great ape's actual life history remains unknown. The goal of this paper is to present mostly new, very carefully vetted data on orangutan life history, specifically immature survival and birth intervals, and to explain the observed differences with the other hominids, particularly chimpanzees. We combine data from seven current long-term research sites with detailed demographic records (over at least 10 years) on regularly encountered females, for two study populations of Sumatran orangutans (*Pongo abelii*: Ketambe and Suaq Balimbing) and five Bornean populations (*Pongo pygmaeus wurmbii*: Gunung Palung, Sabangau, Tuanan, and *P.p. morio*: Danum Valley and Lower Kinabatangan). Despite many years of investment, for each site the sample size for known birth

dates (within a month) remains small due to the inherent challenges of studying a non-gregarious wild primate. By combining our data, we provide the best-available evidence for the orangutans' extremely long IBIs on both islands as well as their extremely low infant mortality compared with all other apes, including most humans. Published data of smaller sample sizes suggested a gradient from west to east of decreasing interbirth intervals, which was interpreted to reflect higher mortality risks due to greater periodic fruit scarcity in (east) Borneo compared to (northwestern) Sumatra (e.g., Wich et al., 2004; van Schaik et al., 2009). Our larger dataset allows us to test whether there are differences in wild orangutan interbirth intervals, infant survival, and female age at first birth and thus in their life history by (sub-) species and geographical range.

Methods

We compiled data from the long-term orangutan field sites for which regular observations over at least 10 years were available (Fig. 1, Table 1). For *P. abelii*, long-term data are available from only two sites west of the Alas river (a genetically distinct population: Nater et al. 2013), in relatively productive forests (Marshall et al., 2009a). For Bornean orangutans, data could be included only for *P.p. wurmbii* from three locations and *P.p. morio* from two sites in Sabah.

For each site we classified births (Table 2) as 'known' when the birth date was known to within a month (based on sightings before and after birth), 'estimated' when birth dates were assessed with a margin of at most three months, 'roughly estimated' when birth dates could be assigned with at least six months accuracy and 'very rough estimates' when very young and dependent infants were assessed by experienced field workers to be less than three years old at

first encounter. IBIs were classified as ‘known’ when both birth dates were ‘known’, as ‘estimated’ when both were at least ‘estimated’, and as ‘roughly estimated’ when both were at least rough estimates. No IBI was assigned if one of the birth dates was unknown. Age at first reproduction was classified by the same criteria. These criteria are stricter and more accurate than those used in previous publications. Unless otherwise noted, analyses were restricted to infants of known sex, to known, estimated and roughly estimated values, and to fully wild individuals only. This final criterion served to exclude some formerly re-introduced individuals and their descendants from analyses of the data on the Ketambe population. Sex ratio among immatures was measured as the proportion of males among the infants with at least a roughly estimated birthdate. Offspring are referred to as dependent (‘infant’) when they are still seen in nipple contact with the mother multiple times per day, presumably receiving at least some milk, and having the opportunity to do this at night by sharing the mother’s night nest (van Noordwijk and van Schaik, 2005; van Noordwijk et al., 2013). Accordingly, we estimated final weaning (end of lactation), which was not systematically monitored in all populations or for all individuals, as the point in time when the immature started to sleep in its own night nest.

Pre-weaning mortality

Death of a known dependent offspring was assumed when the mother was seen for more than a full day without her unweaned offspring, and the immature was not later found to be alive. We measured the proportion of infants that survived at least until the birth of the younger sibling, or at least until the first offspring was completely weaned. However, in all known cases weaning preceded the birth of the next offspring. Thus our measure of pre-weaning mortality is conservative.

Interbirth intervals

These were calculated between live births if the first of these offspring survived at least to weaning. Right-censored IBIs (i.e., ‘open’, unfinished intervals in which no new infant was born yet) were calculated only when the current infant was alive at last sighting of the mother, but she had not yet given birth to her next offspring. The estimates ignore known or suspected pregnancy failures. In several cases females showed positive results on standard (locally available) human pregnancy tests, but failed to give birth to a healthy offspring. In the wild Ketambe population, two actual miscarriages with bloody tissue discharge were witnessed (Utami Atmoko pers. obs.). However, because it is impossible to systematically include such cases in all but extremely closely monitored populations, these data were not included in the analyses. Thus, as in other studies, we report IBIs between known infants only, and similarly, mortality rates for known infants only (e.g., Muller and Wrangham, 2014). Such estimates thus ignore potential perinatal mortality, which could have been missed since individuals were rarely followed every day. When a previous infant had died before weaning, we determined the interval until the next birth after the death of the unweaned offspring.

Maternal stage

For each birth and subsequent birth interval, mothers were classified (Table 2) based on their known reproductive histories and assessments based on visible body condition by the researchers: ‘young’ (when they had been known as adolescents before the birth of their first offspring and/or estimated to be <21 years old and thus having at most 1 living offspring), ‘multiparous’ (>21 years and 2nd- 4th birth), or ‘old’ (birth 5 or later and estimated >35 years). The multiparous age class may include some older females whose older offspring was not recognized as such, had died or had disappeared from the study area. In particular, earlier male offspring of the older mothers may be missed, since in both orangutan species male offspring

disperse before becoming reproductively active (Morrogh-Bernard et al., 2011; Arora et al., 2012; Nietlisbach et al., 2012; van Noordwijk et al., 2012). In addition, when no genetic parentage analyses were available, some earlier daughters may also not have been recognized as such and ignored in the reproductive history of the mother. Likewise, daughters of peripheral females who had settled outside the study area might have been missed. We therefore consider the assignment of females to the ‘old’ class as a minimum estimate.

Statistics

Since short intervals are more likely to be completed than longer ones within the limited duration of a study, the average or median value of completed IBIs and age at first reproduction would be biased in favor of these shorter intervals. Therefore, we conducted survival analyses on all data (including right-censored data). To visualize IBIs, survivorship and age at first reproduction, we used the ‘survival’ package (Therneau, 2015a) for R (version 3.3.0, R Core Team, 2016) to compute and plot Kaplan-Meier survival curves. To assess the probability of an event occurring in each of these three life-history parameters, we fitted Cox proportional hazard mixed-effects models to our observations on each of them, using the ‘coxme’ package (Therneau, 2015b).

For age at first reproduction, our initial, very strict, data-quality criteria left us with too few observations for even the simplest model to converge. Therefore, we report simple summary statistics for this life-history parameter in the Results section. For all descriptive statistics we report averages and standard deviations. IBI and survivorship could successfully be expressed as proportional hazard functions of subspecies (*P. abelii* vs. *P.p. wurmbii* vs. *P.p. morio*), infant sex, and maternal stage, while accounting for the hierarchical structure of our

data by incorporating individual identity nested within study-site, nested within subspecies as random effects in the Cox models. Overall model performance was assessed by Likelihood Ratio Tests (LRT), comparing the fit of fully parameterized models to their respective null models (intercept and random structure only).

Results

Age at first reproduction

Age at first reproduction (AFR) was calculated only if both the birth month of the female herself and the month in which she gave birth to a viable offspring for the first time were estimated with an accuracy of at least six months. Only four AFRs could be calculated based on these criteria, yielding an average of 178.0 ± 17.0 months or 14.8 years ($n=3$ for Sumatra, $n=1$ for Borneo). If we also included very rough estimates, including females whose own birth year was estimated with a margin of > 6 months (estimated when they were small, dependent immatures at first encounter with the mother – as was done in most of the previously published wild cases) AFR ranged from 152 to 196 months with an average of 172.4 ± 12.7 months ($n=13$), with no apparent difference between the islands (Sumatra 172.0 ± 11.9 , $n=5$; Borneo 172.6 ± 13.9 , $n=8$; Welch's t-test: $t_{9.74} = 0.086$, $p = 0.933$, $r = 0.028$).

A Kaplan-Meier survival analysis (Fig. 2) on all data from the seven sites, including 41 censored female ages and the four known AFRs, but excluding the very rough estimates, is consistent with a median age of first reproduction at around 14.5 years. Because sample size remains very small, no further statistics were calculated, but there was no clear difference between populations, subspecies or species (Supplementary Online Material [SOM] Table S1 for a Cox model fitted to the dataset based on less restrictive criteria of inclusion, and overview in

SOM Table S5). Similarly, sample size is still too small to analyze possible other effects such as maternal age on a daughter's AFR.

Offspring sex ratio

Even though in the past the smaller samples from some sites seemed to indicate a biased sex ratio at birth, the current larger sample does not show either an overall bias or a difference between the islands. Among the 103 infants with known or closely estimated birth dates, the sex ratio was slightly male-biased on both islands (for Sumatra 13f, 17m, 1 unknown sex; $n=31$; for Borneo 33f, 36m, 3 unknown; $n=72$; Fischer exact tests on either the 2x3 or 2x2 contingency table – omitting the unknowns - are not significant ($p= 0.928$ and $p= 0.87$, respectively). The sample sizes per site are still too small to test for any finer-grained geographic patterns. Young and old mothers may have more male offspring, but the differences were not significant (Fischer exact tests on either the 2x3 or 2x2 contingency table NS).

Interbirth interval after surviving infant

For the interval between live births in which the first offspring survives until weaning, the Kaplan-Meier curves and Cox proportional hazards models (Fig. 3; Table 3) show no significant differences between the islands, or subspecies (see also SOM Table S4 values per site). The mean overall IBI based on completely known intervals was 91.5 ± 23.7 months or 7.6 ± 2 years ($n=16$), whereas for the larger dataset, which includes estimated and roughly estimated intervals, the mean was 93.7 ± 21.4 months or 7.8 ± 1.8 years ($n=49$). Three females estimated to be over 45 years old, who had successfully weaned their last offspring, apparently did not give birth to a subsequent offspring for at least 132, 168 and 180 months after the previous

birth, respectively. None of the other open intervals with a reliably estimated start date was longer than 80 months and thus were all still within the range of observed closed intervals. The 'tail' in the Sumatran distribution of completed intervals seems to be longer, with three outliers of 12 years or more, suggesting exceptionally long IBIs could be more common there, although we cannot completely rule out early infant loss during this period. However, the average of the remaining Sumatran intervals (89.5 ± 12.6 , $n=20$) was very similar to the Bornean intervals (89.4 ± 13.4 , $n=26$; Welch's t-test: $t_{42,24} = -0.017$, $p = 0.987$, $r = 0.003$). When the outliers are included the difference remains insignificant (Wilcoxon rank-sum test: $W = 340$, $p = 0.417$, $r = -0.116$), and median values for Sumatra and Borneo (90.0 and 89.5 months, respectively, based on 23 vs. 26 values) were almost identical. The birth interval after raising an infant to weaning seemed unaffected by offspring sex, but tended to be shorter for multiparous mothers (NS: Table 3). In some sites, conceptions appear more likely at the onset of irregularly spaced periods with superabundant fruit, leading to clustered births (e.g., Gunung Palung: Knott, 1999, 2001; Knott et al., 2009; Danum Valley: Kuze et al., MS; Lower Kinabatangan: Oram, 2018). This might lead to more extremely short or long intervals, but so far this effect is not visible in the data.

Interval after infant death

There were only eight cases in total to estimate the interval after the death of previous offspring until the birth of the next one (within at least a few months), which yielded an average of 14.0 ± 4.2 months. Thus mothers tended to need approximately half a year to conceive again after the death of their previous, unweaned offspring (see Fig. 4: assuming an 8-months gestation: Sodaro et al., 2007). Because this interval was not affected by the age of the offspring at the time of death (Borneo only: $r_s = -0.31$, NS), time needed to 'recover' did not

appear to be affected by the duration of their investment in the previous offspring or how close that offspring was to weaning.

Offspring survival

To estimate offspring survival, Kaplan-Meier survival analyses were performed including all offspring with known, estimated and roughly estimated birth dates and including those still alive at the end date of the study, or for infants, the last verified sighting of the mother (censored data). Fig. 5a shows the extremely high survivorship of infants with known birth month: 91% (combined males and females) survived to at least weaning age (see below) and even until at least 10 years of age. No deaths were reported for offspring between weaning and 15 years of age, although at least one juvenile, whose birth date could not be estimated reliably, died during the early years of study in Danum Valley (Kanamori et al., 2012). There was no statistically significant difference between Sumatra and Borneo (all sites combined) in pre-weaning survival, but pre-weaning survival in the two *P. p. morio* sites sampled in this study may have been lower (Fig. 5b; Table 4, SOM Table S5: subspecies difference). Overall there was a tendency for males to have slightly lower survival (Fig 5c) up to age 10. Since males tend to roam widely and eventually disperse, starting at c. 10-15 years of age, the survival data beyond 10 years are more reliable for females. In our combined sample, offspring of multiparous mothers have a higher survivorship than those of young and old mothers (see methods for maternal age class criteria; Fig. 5d). However, note that survival up to 10 years for offspring of old mothers is still around 80%.

In most cases, the cause of death of an infant could not be determined (SOM Table S2). However, one early infant death (7-10 days) in the Lower Kinabatangan (Oram et al.,

unpublished) was attributed to pulmonary edema. This infant had rapidly weakened between the initial sighting at 1-3 days of age to the second sighting at 6-9 days of age. This infant's experienced mother (this was her 5th known offspring – the older ones were genetically confirmed) seemed to be aware that something was not right as she spent a lot of time inspecting and repositioning the infant to try to help it nurse in ways she had not been observed doing with previous offspring, milk was noted to repeatedly leak out of the infant's mouth suggesting it could not swallow properly. This female subsequently gave birth to a healthy infant the following year.

In Danum Valley, a small but weaned immature (estimated >3 years old at first encounter – not included in current analyses) was found severely wounded a week before she succumbed to wound infection (Kanamori et al., 2012). This immature was already ranging alone after the birth of a younger sibling. The fatal wounds were consistent with an attack by a Sunda clouded leopard, *Neofelis diardi*, although it had not succeeded in killing the immature outright, despite the latter's relatively small size. Another apparent predator attack in Tuanan resulted in fatal injury to the mother but left the infant unharmed (SOM Table S2). In addition, in Kinabatangan two orangutan mothers, whose infants were playing nearby, have been observed to mob and chase a clouded leopard they had detected in the canopy (Oram, unpublished). Male infanticide was suspected in one infant loss in Gunung Palung (Knott et al., unpublished), but no other (suspected) cases have been reported in the wild.

Comparative survival

Since ages of most adult females are not known, we cannot yet construct a complete life table. However, in all sites adult female deaths are very rare. For the Tuanan resident females (>15 years of age), for instance, only two, both estimated to be >35 years old, have disappeared (and were confirmed dead: Marzec et al., 2016) in 146 adult female years, for an average yearly mortality rate of < 1.5%.

Our data from the seven sites do allow for a comparison of female survival from birth up to 15-18 years of age with data on captive orangutans (Wich et al., 2009a), wild chimpanzees (Kanyawara: Muller and Wrangham, 2014; Ngogo: Wood et al., 2017), wild gorillas (Bronokowski et al., 2016b), and traditional human societies (Hadza: Blurton-Jones 2016; combined survival up to 15 years in 13 small-scale human societies: Walker et al., 2006) (Fig. 6; Table 5). We also added the female survival data of the current human female population of Switzerland (Human Life-Table Database, <http://www.lifetable.de>; see also SOM Table S3). Actually, the Swiss girls fall within the 95%-CI of wild orangutan females, so although survival of the former is higher, this difference is not statistically significant given the uncertainty around our estimate for wild orangutans. To broaden the perspective, we also included the survival data for female Amboseli elephants (Lee et al., 2011), which have a similarly slow life history, with IBIs (4-5 years) in the same range as chimpanzees (Moss and Lee, 2011). Even though early elephant survival rates are higher than all chimpanzees and hunter-gatherer populations (Fig. 6), they still have lower immature survival values than wild orangutans in our sample.

Discussion

Orangutan life history

Despite several decades of field research, only a few orangutan females are known from birth to their AFR. The average AFR of 14.8 years for the four known cases is only half a year higher than when an additional nine females, with (mostly very roughly) estimated ages, are included: 14.3 years. Thus, earlier estimates of AFR at 14-15 years of age (Knott, 2001) were confirmed in our combined dataset. Studies covering at least 20 consecutive years will be needed to test whether there are systematic differences between populations, or what conditions affect relatively early or late AFRs.

Our dataset on IBIs with a median of 7.5 years confirms the long mean interbirth interval reported based on earlier, smaller samples (Galdikas and Wood, 1990; Knott, 2001; Wich et al., 2004). Pregnancy failure affects IBIs, and we recorded multiple cases of suspected pregnancy, based on human pregnancy tests and/or labial swelling (which appear within weeks of conception: Sodaro et al., 2007). For example, in 20% (three of 15) of the suspected pregnancies in Tuanan, the female was later seen before the expected due date but no longer pregnant and receiving renewed male sexual interest. In addition, two miscarriages of bloody tissue without the shape of a full-term infant were witnessed in Ketambe, and in most other sites pregnancy failure was suspected at least once. Early pregnancy loss is a common phenomenon in humans as well (10-20% after the first month of gestation: Avalos et al., 2012) and often related to fetal abnormalities. The frequency in orangutans appears to be in the same range, but early fetal loss and later miscarriage are hard to document under field conditions. There is little indication that pregnancy failures were due to poor condition (cf. Brockman and van Schaik, 2005). For example, in Tuanan, all three females known to be pregnant before the onset of the long period of Kalimantan-wide haze due to fires in 2015, subsequently gave birth to healthy infants.

Interbirth interval estimates are systematically overestimated when early infant mortality is missed. Since many females in this dataset were seen irregularly and certainly not on a monthly basis, this possibility must be examined. Reassuringly, among the known births the calculated mortality during the first three months was very low (only two in 66 known births and 0 in 38 births with estimated birth dates and 1-3 months of age), suggesting early mortality is quite low, and thus our estimates are largely unbiased. Hassell et al. (2017) found that in wild mountain gorillas early infant mortality attributable to fetal and developmental problems is around 10% of all mortality up to 3.5 years. However, even if we assume a similar value for early mortality in orangutans, i.e., if we had missed an extra 10% of early infant mortality in our field data because observations were intermittent, the infant survival rate would still be considerably higher in the orangutan sample than reported for any other apes or pre-industrial human populations.

In our compilation of data on life history parameters we found not only a very high infant survival rate (94% for female and 87% for male infants up to weaning), but we also did not record any mortality among females known since weaning until AFR across all sites, and only one instance in the larger sample of known individuals. Similarly, males have high survival rates during their early post-weaning years, after which their disappearance from the study sites may be due to dispersal rather than death. We did not find evidence for increased mortality around or following weaning (as in some chimpanzee populations: Hill et al., 2001), suggesting offspring have sufficient ecological competence at the time of weaning (van Noordwijk and van Schaik, 2005; Schuppli et al., 2016).

Earlier studies suggested (west-) Sumatran orangutans (*P. abelii*) have an even slower life history than the Bornean orangutans (*P. pygmaeus*), with the population of *P. p. morio*

having the fastest life history (Wich et al., 2004; van Schaik et al., 2009). In our larger dataset, although still small compared to other primate species, we found no evidence for systematic differences in interbirth intervals between the study populations. This was true for all other life history parameters as well, with the possible exception of slightly higher infant mortality in *P. p morio*. This lack of life history differences between Sumatran and Bornean orangutans was also reported in studies of captive orangutans that either found no inter-island difference in IBI, AFR or infant mortality (Cocks et al., 2007a,b; Anderson et al., 2008; Wich et al., 2009a) or the opposite of the early wild data (Anderson et al., 2008).

Adult female life tables cannot be constructed yet, but available evidence (Wich et al. 2004; Marshall et al., 2009b; Tuanan data) suggests adult female yearly mortality rate during most of the lifespan is < 2%. Longevity remains unknown, but is likely to be more than 50 years, although very old females appear to range less widely and become more solitary, at least at Tuanan (van Noordwijk et al., unpublished). Indeed, we found that middle-aged (multiparous) females tended to have shorter interbirth intervals and had higher infant survival than either primiparous or old females. This pattern mirrors that found in many other species (van Noordwijk & van Schaik, 1999; Bronikowski et al., 2016a; Lee et al., 2016).

In sum, current evidence suggests an age at first reproduction of nearly 15 years, infant survival until weaning of around 90% and a mean IBI of around 7.6 years for all wild orangutans. Clearly, more detailed comparisons at the population level are warranted, but we should not expect these taxon-level averages to change much in the future.

Causes of mortality

All unweaned offspring that disappeared were assumed dead, but the cause was usually not established (SOM Table S2) because bodies were rarely found. In most cases, the infant looked healthy when last sighted shortly before disappearance. Since mothers usually conceived again within half a year from when their pre-weaned offspring had died, malnourishment or poor maternal condition was an unlikely cause in most cases reported here. Respiratory or other contagious diseases are often associated with both infant and adult mortality in the more gregarious and terrestrial chimpanzees (Kuehl et al., 2008; Wood et al., 2017) and gorillas (Hassell et al., 2017), but there is no evidence for this in wild orangutans.

Predation, though rare, is not absent (see SOM Table S2). The observed case of mobbing of a clouded leopard in Kinabatangan (Oram, unpublished) was very similar to witnessed mobbing by other arboreal primates (e.g., gibbons and langurs: Wilcox et al., 2017). This suggests that clouded leopards are an important threat for immatures; indeed, they are known to kill orangutans in rescue sites (Rijksen 1978; Phillips and Phillips 2016). Even though male infanticide is a frequent cause of early death in chimpanzees and gorillas (overview in Palombit, 2012; Hassell et al., 2017), we know of only one suspected case of infanticide in wild orangutans (Knott et al., unpublished) and one known case in captivity (Mallinson et al., 1984). Although it is generally assumed that male orangutans, especially in Borneo will not consistently benefit from infanticide (van Schaik, 2004; Beaudrot et al., 2009), females were found to conceive within six months after the loss of their infant on average (Fig. 4), so males might derive a fitness benefit from killing infants.

Unlike in chimpanzees and gorillas (Hassell et al., 2017; Wood et al., 2017), by elimination, we suggest that the most likely cause of death for pre-weaning orangutans may well be accidents, such as falls from the canopy either during social interactions, solitary play or

travel. Although such falls were frequently observed, we don't have hard evidence so far on fatal outcomes. Similarly, in our comparison with the species with the next lowest infant mortality, African elephants, 'accidents' are recorded as a major cause of death of young calves, in addition to mortality due to poor nutrition and predation (Moss and Lee, 2011), whereas disease does not seem to play a major role. In this long-lived species, living in matrilineal herds, the protection provided by maternal relatives contributes to the calves' early high survival.

In conclusion, wild orangutans appear to have largely eliminated contagious disease and starvation, and strongly reduced predation and, probably, infanticide as causes of immature mortality, leaving accidents as the most common, albeit still rare, cause.

Plasticity versus maladaptation

Studies of reproductive ecology suggest that in primates, including humans and chimpanzees, higher energy availability leads to earlier AFR and shorter IBIs (Ellison et al., 1993; Knott, 2001; Emery Thompson, 2013). Orangutans likewise speed up growth and reproduction when food is more reliably available (Knott et al., 2009). Provisioned rehabilitant orangutans on both islands are reported to give birth several years earlier than wild ones (Galdikas and Ashbury, 2012; Kuze et al., 2012). Mothers in captivity tend to have IBIs two years shorter (Cocks, 2007a, b; Anderson et al., 2008) than the median IBI of 7.5 years (this study) found for wild orangutans, even when the previous offspring was housed continuously with its mother (Shumaker et al., 2008). Both earlier AFR and shorter IBIs suggest a strong effect of physical condition and health, and possibly social (and management) conditions, on these reproductive parameters (Knott, 2001). This suggests high plasticity in the rate of development and reproduction in orangutans.

Among mothers in rehabilitation centers, infant mortality (at 18-61%) was found to be much higher than in the wild, which was attributed to poor mothering skills, increased rates of disease transmission in high-density release sites with fixed feeding locations (Kuze et al., 2012), and high parasite prevalence accompanying high use of the ground around the feeding sites (Foitova et al., 2009). However, infant mortality rate in zoo-housed orangutans is similarly high (Cocks, 2007b). Thus, despite nutritional advantages, veterinary care and freedom from predation in captivity and rescue centers, infant mortality is much lower in the wild (Fig. 6).

Although mean survivorship has generally increased significantly for orangutans in captivity since 1965, and some individuals are known to be over 50 years old, median survival of captive-born maternally raised Sumatran (*P. abelii*) females was calculated to be only 24.6 years (based on international studbook data since 1986: Wich et al., 2009b). In another comprehensive captive study, the age at last reproduction was found to be 41 years (Shumaker et al., 2008). To date, good longevity and life time reproductive records of wild orangutans are still scarce, but in all populations under long-term observation in this study, females in their mid-to-late 30s are well documented and at least some females are known to be reproducing well into their mid-to-late 40s.

The discrepancy between reproductive parameters, such as AFR and IBI, under natural vs. 'human-enhanced' conditions suggests a shift in resource allocation when an individual grows up without experiencing scarcity. However, there are indications that plasticity allowing accelerated maturation (early AFR) and increased reproductive rate (shorter IBIs) involves a life-history tradeoff. The relatively high infant mortality reported in rescue centers and zoos holds especially for second-generation primiparous mothers with very early AFR (< 12 years old) (Cocks, 2007b; Galdikas and Ashbury, 2012). Moreover, zoo-housed females with shorter IBIs

tend to die at a younger age (Cocks, 2007b). All this suggests that the observed response to superabundant food conditions is actually maladaptive. Apparently, such conditions are so far outside the natural range in extent and duration that natural selection did not need to cap the open-ended response to improved conditions. Similar negative consequences of deviations from the average life history pattern have also been reported for other long-lived species with slow life history (e.g., red deer: Nussey et al., 2006, elephants: Hayward et al., 2014; and humans: King, 2003; Hayward et al., 2015). This finding has obvious and important implications for captive management.

Orangutans and chimpanzees compared

The new results on both orangutans and chimpanzees (Muller and Wrangham 2014; Wood et al., 2017) suggest similar life-history variables, but with a consistent difference: orangutans are slower. Orangutan IBIs are longer (7.5 years versus 5-6 years: Watts, 2012), indicating higher reproductive effort. AFR is later at 14.8 years, especially when compared with the chimpanzee females that breed in their natal community (c. 13.5 years; Walker et al., 2018). Although this is not the modal situation in chimpanzees, it is actually the more meaningful comparison of growth and development with the female-philopatric orangutan because delays due to dispersal do not reflect unconstrained energetic allocation. Finally, the very high adult female survival in orangutans (this study, see also Wich et al., 2004) relative to most chimpanzee populations (Muller and Wrangham, 2014; Wood et al., 2017) is consistent with delayed senescence in orangutans. Thus, orangutans are simply slower than chimpanzees.

This slowdown in orangutan life history is consistent with the rarity of observed disease or predation in wild populations, since it suggests that conservative growth, development and

reproduction could actually pay off in improved survival. The very conservative reproductive investment may reflect the need to deal with fluctuating food abundance (Knott, 2001), including multiple periods of food scarcity during the years of extended lactational effort (van Noordwijk et al., 2013). The arboreal lifestyle limits opportunities for fat storage to buffer fluctuations in food intake (Heldstab et al., 2017), which further contributes to the risk of maternal energy depletion in orangutans (Knott, 2001, Knott et al., 2009). The slowdown also enables the hypometabolism found in orangutans (Pontzer et al., 2016). The extreme conservatism in growth may explain the remarkable near-absence of any starvation-induced mortality, even during extended periods with extremely low food abundance. If conceptions are more likely during increasing fruit abundance, one may expect a strong influence of food scarcity on maternal condition and thus infant survival. Yet, mortality rates were equally low at sites that experience irregular mast-fruiting and sites with less extreme variation and higher fruit abundance, e.g., Suaq Balimbing. In addition, even during the extended period of fruit scarcity in Tuanan following the fires and Kalimantan-wide severe haze for c. three months in 2015 (unpublished data), no known immatures were lost and three infants born during the post-haze scarcity period all survived at least their first year of life.

Overall, then, the chimpanzee is the faster ape because it is more terrestrial and more gregarious. The orangutan's extreme arboreality and low gregariousness protects it from the threats of predation and spread of contagious diseases that affect the more terrestrial hominids. Because terrestriality is obviously a derived trait in great apes, given its morphology adapted to below-branch arboreal locomotion (Thorpe and Crompton, 2006), the orangutan may actually reflect the ancestral great ape state. This speculation is supported by the

remarkably slow life history of the relatively very small, but exclusively arboreal gibbons, with IBIs of c. 3.5 years and an AFR of 11 years (Reichard and Barelli, 2008), in the gorilla range.

Implications for human evolution

Orangutan life history illustrates how the basic hominoid bauplan can give rise to extremely slow life histories when conditions enable very low unavoidable mortality. Paradoxically, low habitat productivity may have enforced a near-solitary lifestyle, which in combination with arboreality, allows orangutans to largely avoid both predation and contagious diseases. In contrast, most likely, modern humans only reached similar or higher survival probabilities than female orangutans in the 20th century (SOM Table S3). Even though pre-contact hunter gatherer populations may have suffered much lower mortality due to disease (Gurven and Kaplan, 2007), all current populations represented in Fig. 6 fall outside the confidence limits for orangutans from c. six months of age onward.

Although ape longevity remains only crudely estimated, we can now see why humans differ from the basic hominoid pattern. First, extensive allomaternal care allowed for shortened IBIs (Knott 2001; Robson et al., 2006; Kramer and Ellison, 2010; van Schaik, 2016). Second, weaponry and cooperative defense reduced unavoidable mortality due to predation, despite a fully terrestrial lifestyle, allowing longevity to be maximized (Isler and van Schaik, 2012). Third, very low densities (Kelly, 2013) ensured low mortality due to infectious disease (Black, 1975). Finally, development slowed down due to the strong increase in brain size and the resulting competition for energy during development (Isler and van Schaik, 2012; Kuzawa et al., 2014).

Concluding remarks

Before getting overly optimistic about the orangutans' ability to survive in low-productivity forest thanks to their extremely slow life history, it should be remembered that our long-term study sites are situated in some of the best remaining larger forest areas, despite a selective logging history in some sites, and border areas with increasing disturbance for most sites. Orangutans living in smaller remnant forest patches are unlikely to achieve these high survival rates found in larger and rather undisturbed forests. Their exceptionally slow life history leaves orangutans vulnerable to unfettered anthropogenic change (such as hunting and habitat loss due to land conversion; Marshall et al., 2009b), which may affect health and infant and maternal survival. Such effects could result in catastrophic population crashes, but the ability of adults to endure periods of starvation (perhaps especially in Borneo: Vogel et al., 2012) may mean these crashes are not numerically visible until it is too late.

The data presented here are still modest, especially with respect to longevity. The big differences in reproductive timing between wild and captive orangutans illustrate that it is important to collect long-term data on natural populations to gain an understanding of their life history (Borries et al., 2013). This should further increase our motivation to protect and enable healthy populations of great apes to survive in their natural habitats.

Funding

For major funding at Ketambe, Suaq Balimbing, and Tuanan we thank NWO/WOTRO, WCS, L.S.B. Leakey Foundation, A.H. Schultz Stiftung, University of Zurich, Utrecht University, Paneco, Swiss National Science Foundation and Philadelphia Zoo. Research at Gunung Palung was supported by grants to CDK from the National Science Foundation (BCS-1638823BCS, BCS-0936199, 9414388); the National Geographic Society; the L.S.B. Leakey Foundation; the Wenner

Gren Foundation; the US Fish and Wildlife Service (F15AP00812, F12AP00369, 98210-8-G661); Conservation, Food and Health Foundation; Disney Wildlife Conservation Fund; and Nacey Maggioncalda Foundation. For work in Danum Valley NK thanks JSPS Core-to-Core Program, A.Advanced Research Networks "Tropical Biodiversity Conservation" Wildlife Research Center, Kyoto University, the Leading Graduate Program in Primatology and Wildlife Science, JSPS Research Fellowships for Young Scientists (21-634, 25-40012, 29-40025). FO thanks long-term support to HUTAN–KOCF by Arcus Foundation; zoos of Zooparc de Beauval, la Palmyre, Chester, Woodland Park, Houston, Cleveland, Columbus, Phoenix, Saint Louis, Basel, Apenheul, Hogle, and Oregon Metroparks; Association of Zoos and Aquariums Great Ape Taxon Advisory Group; Australian Project; Synchronicity Earth; United States Fish and Wildlife Service; World Land Trust; Waterloo Foundation; and other partners. HM-B thanks support for Sabangau from Wildlife Conservation Society (WCS), the US Fish and Wildlife Service Great Ape Conservation Fund, Borneo Nature Foundation, Primate Conservation Inc., the L.S.B. Leakey Foundation and Arcus.

Acknowledgements

For allowing long-term research efforts in multiple locations, granting permissions to many subsequent local and foreign students and researchers, we're very grateful to the government agencies in Indonesia (in particular LIPI, RISTEKDIKTI, Ministry of the Interior, Ministry of Environment and Forestry, KPH, BKSDA, Gunung Leuser NP, Gunung Palung NP, Sabangau NP), and in Malaysia (Economic Planning Unit, Sabah Biodiversity Center, Danum Valley Management Committee, Sabah Wildlife Department). We thank local NGOs and

598 universities for their sponsorship and long-term collaboration: Universitas Nasional (UNAS),
599 Universitas Tanjungpura (UNTAN), Universitas Palangkaraya (UNPAR, in particular former rektor
600 Napa J. Awat) and the late Dr. Suwido Limin, Director of CIMTROP (Centre of International Co-
601 operation in Management of Tropical Peatlands), Universiti Malaysia Sabah, the Gunung Palung
602 Orangutan Conservation Program/Yayasan Palung, the Eijkman Institute for Molecular Biology,
603 the Center for Research and Development in Biology (PPPB), Paneco/YEL, BOSFoundation, BOS-
604 Mawas Conservation program, BNF, Borneo Nature Tours SDN, BHD.

605 We are indebted to our many colleagues as well as many field managers, all field
606 assistants and students who contributed their dedication and observations over many decades.
607 In particular we thank Tatang M. Setia, Abu Hanifah Lubis, Fitriah Basalamah, Serge Wich, Erin
608 Vogel, Lynda Dunkel, Brigitte Spillmann, Anna Marzec, Julia Kunz, Ellen Meulman, Amy Scott,
609 Katherine Scott, Tri Wahyu Susanto, Mark Harrison, Rebecca Armson, Jennifer Brousseau,
610 Sophie Kirlin, Tomoko Kanamori, Saika Yamazaki, Renata Mendonça, Hamisah Elahan, Rusiman
611 Rukimin, Adul Rajak Sharon, Azli Etin, Herman Suali, Waslee Maharan, Mohd. Daisah Kapar,
612 Hartiman Abd. Rahman and Mohd. Faisal Asmara for assistance with compiling these data.
613 Finally we thank Anne Russon, Alison Ashbury, Matt Novak, and Erin Vogel for discussion, and
614 the anonymous reviewers for their helpful feedback.

References

- Anderson, H.B., Emery Thompson, M., Knott, C.D., Perkins, L., 2008. Fertility and mortality patterns of captive Bornean and Sumatran orangutans: is there a species difference in life history? *Journal of Human Evolution* 54, 34-42.
- Arora, N., van Noordwijk, M.A., Ackermann, C., Willems, E.P., Nater, A., Greminger, M., Nietlisbach, P., Dunkel, L.P., Utami Atmoko, S.S., Pamungkas, J., Perwitasari-Farajallah, D., van Schaik, C.P., Krützen, M., 2012. Parentage-based pedigree reconstruction reveals female matrilineal clusters and male-biased dispersal in nongregarious Asian great apes, the Bornean orang-utans (*Pongo pygmaeus*). *Molecular Ecology* 21, 3352-3362.
- Ashton, P.S., Givnish, T.J., Appanah, S., 1988. Staggered flowering in the Dipterocarpaceae: new insights into floral induction and the evolution of mast fruiting in the aseasonal tropics. *American Naturalist* 132, 44-66.
- Avalos, L.A., Galindo, C., Li, D.-K., 2012. A systematic review to calculate background miscarriage rates using life table analysis. *Birth Defects Research (Part A)* 94, 417-423.
- Beaudrot, L.H., Kahlenberg, S.M., Marshall, A.J., 2009. Why male orangutans do not kill infants. *Behavioral Ecology and Sociobiology* 63, 1549-1562.
- Black, F.L., 1975. Infectious disease in primitive societies. *Science* 187, 515-518.
- Blurton Jones, N., 1986. Bushman birth spacing: A test for optimal interbirth intervals. *Ethology and Sociobiology* 7, 91-105.
- Blurton Jones, N., 2016. *Demography and Evolutionary Ecology of Hadza Hunter-Gatherers*. Cambridge University Press, Cambridge.
- Borries, C., Gordon, A.D., Koenig, A., 2013. Beware of primate life history data: A plea for data standards and a repository. *PLoS ONE* 8, e67200.

- 638 Brockman, D.K., van Schaik, C.P., 2005. Seasonality and reproductive function. In: Brockman,
639 D.K., van Schaik, C.P. (Eds.), Seasonality in Primates: Studies of Living and Extinct Human
640 and Non-Human Primates. Cambridge University Press, Cambridge, pp. 269-305.
- 641 Bronikowski, A.M., Cords, M., Alberts, S.C., Altmann, J., Brockman, D.K., Fedigan, L.M., Pusey, A.,
642 Stoinski, T., Strier, K.B., Morris, W.F., 2016a. Female and male life tables for seven wild
643 primate species. Scientific Data 3, 160006.
- 644 Bronikowski, A.M., Cords, M., Alberts, S.C., Altmann, J., Brockman, D.K., Fedigan, L.M., Pusey, A.,
645 Stoinski, T., Strier, K.B., Morris, W.F., 2016b. Data from: Female and male life tables for
646 seven wild primate species. Dryad Digital Repository.
647 <https://doi.org/10.5061/dryad.v28t5>
- 648 Cocks, L., 2007a. Factors affecting mortality, fertility, and well-being in relation to species
649 differences in captive orangutans. International Journal of Primatology 28, 421-428.
- 650 Cocks, L., 2007b. Factors Influencing the well-being and longevity of captive female orangutans.
651 International Journal of Primatology 28, 429-440.
- 652 Ellison, P., Panter-Brick, C., Lipson, S., O'Rourke, M., 1993. The ecological context of human
653 ovarian function. Human Reproduction 8, 2248.
- 654 Emery Thompson, M., 2013. Reproductive ecology of female chimpanzees. American Journal of
655 Primatology 75, 222-237.
- 656 Foitova, I., Huffman, M.A., Wisnu, N., Olsansky, M., 2009. Parasites and their impacts on
657 orangutan health. In: Wich, S.A., Utami Atmoko, S.S., Mitra Setia, T., van Schaik, C.P.
658 (Eds.), Orangutans: Geographic Variation in Behavioral Ecology and Conservation. Oxford
659 University Press, Oxford, pp. 157-169.

- 660 Galdikas, B.M.F., Ashbury, A.M., 2012. Reproductive parameters of female orangutans (*Pongo*
 661 *pygmaeus wurmbii*) 1971–2011, a 40-year study at Tanjung Puting National Park, Central
 662 Kalimantan, Indonesia. *Primates* 54, 61-72.
- 663 Galdikas, B.M.F., Wood, J.W., 1990. Birth spacing patterns in humans and apes. *American*
 664 *Journal of Physical Anthropology* 83, 185-191.
- 665 Gurven, M., Kaplan, H., 2007. Longevity among hunter-gatherers: A cross-cultural examination.
 666 *Population and Development Review* 33, 320-365.
- 667 Halsey, L.G., Coward, S.R.L., Thorpe, S.K.S., 2016. Bridging the gap: parkour athletes provide new
 668 insights into locomotion energetics of arboreal apes. *Biology Letters* 12,
 669 DOI10.1098/rsbl.2016.0608
- 670 Hanya, G., Stevenson, P., van Noordwijk, M., Te Wong, S., Kanamori, T., Kuze, N., Aiba, S.,
 671 Chapman, C. A., van Schaik, C., 2011. Seasonality in fruit availability affects frugivorous
 672 primate biomass and species richness. *Ecography* 34, 1009-1017.
- 673 Hassell, J.M., Zimmerman, D., Cranfield, M.R., Gilardi, K., Mudakikwa, A., Ramer, J.,
 674 Nyirakaragire, E., Lowenstine, L.J., 2017. Morbidity and mortality in infant mountain
 675 gorillas (*Gorilla beringei beringei*): A 46-year retrospective review. *American Journal of*
 676 *Primatology* <https://doi.org/10.1002/ajp.22686>
- 677 Hayward, A.D., Mar, K.U., Lahdenperä, M., Lummaa, V., 2014. Early reproductive investment,
 678 senescence and lifetime reproductive success in female Asian elephants. *Journal of*
 679 *Evolutionary Biology* 27, 772-783.
- 680 Hayward, A.D., Nenko, I., Lummaa, V., 2015. Early life reproduction is associated with increased
 681 mortality but enhanced lifetime reproductive success in pre-industrial humans.
 682 *Proceedings of the Royal Society B: Biological Sciences* 828, 20143053.

- 683 Heldstab, S.A., van Schaik, C.P., Isler, K., 2017. Getting fat or getting help? How female mammals
684 cope with energetic constraints on reproduction. *Frontiers in Zoology* 14, 29.
- 685 Hill, K., Boesch, C., Goodall, J., Pusey, A., Williams, J., Wrangham, R., 2001. Mortality rates
686 among wild chimpanzees. *Journal of Human Evolution* 40, 437-450.
- 687 Isler, K., van Schaik, C.P., 2012. How our ancestors broke through the gray ceiling: comparative
688 evidence for cooperative breeding in early *Homo*. *Current Anthropology* 53, S453-S465.
- 689 Janson, C.H., van Schaik, C.P., 1993. Ecological risk aversion in juvenile primates: slow and
690 steady wins the race. In: Pereira, M.E., Fairbanks, L.A. (Eds.), *Juvenile Primates. Life*
691 *History, Development, and Behavior*. Oxford University Press, New York, pp. 57-74.
- 692 Kanamori, T., Kuze, N., Bernard, H., Malim, T.P., Kohshim, S., 2012. Fatality of a wild Bornean
693 orangutan (*Pongo pygmaeus morio*): behavior and death of a wounded juvenile in
694 Danum Valley, North Borneo. *Primates* 53, 221-226.
- 695 Kelly, R.L., 2013. *The Lifeways of Hunter-Gatherers. The Foraging Spectrum*. Cambridge University Press,
696 Cambridge.
- 697 King, J.C., 2003. The risk of maternal nutritional depletion and poor outcomes increases in early
698 or closely spaced pregnancies. *Journal of Nutrition* 133, 1732S-1736S.
- 699 Knott, C.D., 1998. Changes in orangutan caloric intake, energy balance, and ketones in response
700 to fluctuating fruit availability. *International Journal of Primatology* 19, 1061-1079.
- 701 Knott, C., 1999. Orangutan behavior and ecology. In: Dolhinow, P., Fuentes, A. (Eds.), *The*
702 *Nonhuman Primates*. Mayfield Publishing Company, Mountain View, CA, pp. 50-57.
- 703 Knott, C.D., 2001. Female reproductive ecology of the apes. In: Ellison, P.T. (Ed.), *Reproductive*
704 *Ecology and Human Evolution*. Aldine De Gruyter, New York, pp. 429-463.

- 705 Knott, C.D., 2004. Orangutans: the largest canopy dwellers. In: Lowman, M.D., Rinker, H.B.
706 (Eds.), *Forest Canopies*. Academic Press, San Diego, pp. 322-324.
- 707 Knott, C.D., Emery Thompson, M., Wich, S.A., 2009. The ecology of female reproduction in wild
708 orangutans. In: Wich, S.A., Utami Atmoko, S.S., Mitra Setia, T., van Schaik, C.P. (Eds.),
709 *Orangutans: Geographic Variation in Behavioral Ecology and Conservation*. Oxford
710 University Press, Oxford, pp. 171-188.
- 711 Kramer, P.A., 1998. The costs of human locomotion: maternal investment in child transport.
712 *American Journal of Physical Anthropology* 107, 71-85.
- 713 Kramer, K.L., Ellison, P T., 2010. Pooled energy budgets: resituating human energy allocation trade-offs.
714 *Evolutionary Anthropology* 19, 136-147.
- 715 Kuehl, H.S., Elzner, C., Moebius, Y., Boesch, C., Walsh, P.D., 2008. The price of play: Self-
716 organized infant mortality cycles in chimpanzees. *PLoS One* 3, e2440.
- 717 Kuzawa, C.W., Chugani, H.T., Grossman, L.I., Lipovich, L., Muzik, O., Hof, P.R., Wildman, D.E.,
718 Sherwood, C.C., Leonard, W.R., Lange, N., 2014. Metabolic costs and evolutionary
719 implications of human brain development. *Proceedings of the National Academy of*
720 *Sciences USA* 111, 13010-13015.
- 721 Kuze, N., Dellatore, D., Banes, G.L., Pratje, P., Tajima, T., Russon, A.E., 2012. Factors affecting
722 reproduction in rehabilitant female orangutans: young age at first birth and short inter-
723 birth interval. *Primates* 53, 181-192.
- 724 Lee, P.C., Fishlock, V., Webber, C.E., Moss, C.J., 2016. The reproductive advantages of a long life:
725 longevity and senescence in wild female African elephants. *Behavioral Ecology and*
726 *Sociobiology* 70, 337-345.

- 727 Lee, P.C., Lindsay, W.K., Moss, C.J., 2011. Ecological patterns of variability in demographic rates.
 728 In: Moss, C.J., Croze, H., Lee, P.C. (Eds.), *The Amboseli Elephants. Long-Term Perspective*
 729 *on a Long-Lived Mammal*. The University of Chicago Press, Chicago, pp. 74-88.
- 730 Mallinson, J., 1984. Breeding of great apes at the Jersey Wildlife Preservation Trust and a look
 731 into the future. *Zoo Biology* 3, 1-11.
- 732 Marshall, A.J., Ancrenaz, M., Brearley, F.Q., Frederiksson, G.B., Ghaffar, N., Heydon, M., Husson,
 733 S.J., Leighton, M., McConkey, K.R., Morrogh-Bernard, H.C., Proctor, J., van Schaik, C.P.,
 734 Yeager, C.P., Wich, S.A., 2009a. The effects of forest phenology and floristics on
 735 populations of Bornean and Sumatran orangutans. In: Wich, S.A., Utami Atmoko, S.S.,
 736 Mitra Setia, T., van Schaik, C.P. (Eds.), *Orangutans: Geographic Variation in Behavioral*
 737 *Ecology and Conservation*. Oxford University Press, Oxford, pp. 97-117.
- 738 Marshall, A.J., Lacy, R., Ancrenaz, M., Byers, O., Husson, S.J., Leighton, M., Meijaard, E., Rosen,
 739 N., Singleton, I., Stephens, S., Traylor-Holzer, K., Utami Atmoko, S.S., van Schaik, C.P.,
 740 Wich, S.A., 2009b. Orangutan population biology, life history, and conservation. In: Wich,
 741 S.A., Utami Atmoko, S.S., Mitra Setia, T., van Schaik, C.P. (Eds.), *Orangutans: Geographic*
 742 *Variation in Behavioral Ecology and Conservation*. Oxford University Press, Oxford, pp.
 743 311-326.
- 744 Marzec, M.A., Kunz, J.A., Falkner, S., Utami Atmoko, S.S., Alavi, S.E., Moldawer, A.M., Vogel, E.R.,
 745 Schuppli, C., van Schaik, C.P., van Noordwijk, M.A., 2016. The dark side of the red ape:
 746 male-mediated lethal female competition in Bornean orangutans. *Behavioral Ecology*
 747 *and Sociobiology* 70, 459-466.

- 748 Morrogh-Bernard, H.C., Morf, N., Chivers, D.J., Krützen, M., 2011. Dispersal patterns of orang-
 749 utans (*Pongo* spp.) in a Bornean peat-swamp forest. *International Journal of Primatology*
 750 32, 362-376.
- 751 Moss, C.J., Lee, P.C., 2011. Female reproductive strategies: individual life histories. In: Moss, C.J.,
 752 Croze, H., Lee, P.C. (Eds.), *The Amboseli Elephants: A Long-Term Perspective on a Long-*
 753 *Lived Mammal*. University of Chicago Press, Chicago, pp. 187-204.
- 754 Mul, I.F., Paembonan, W., Singleton, I., Wich, S.A., van Bolhuis, H.G., 2007. Intestinal parasites
 755 of free-ranging, semicaptive, and captive *Pongo abelii* in Sumatra, Indonesia.
 756 *International Journal of Primatology* 28, 407-420.
- 757 Muller, M.N., Wrangham, R.W., 2014. Mortality rates among Kanyawara chimpanzees. *Journal*
 758 *of Human Evolution* 66, 107-114.
- 759 Nater, A., Arora, N., Greminger, M.P., van Schaik, C.P., Singleton, I., Wich, S.A., Fredriksson, G.,
 760 Perwitasari-Farajallah, D., Pamungkas, J., Krützen, M., 2013. Marked population
 761 structure and recent migration in the Critically Endangered Sumatran orangutan (*Pongo*
 762 *abelii*). *Journal of Heredity* 104, 2–13.
- 763 Nietlisbach, P., Arora, N., Nater, A., Goossens, B., van Schaik, C.P., Krützen, M., 2012. Heavily
 764 male-biased long-distance dispersal of orang-utans (genus: *Pongo*), as revealed by Y-
 765 chromosomal and mitochondrial genetic markers. *Molecular Ecology* 21, 3173-3186.
- 766 Nussey, D.H., Kruuk, L.E.B., Donald, A., Fowlie, M., Clutton-Brock, T.H., 2006. The rate of
 767 senescence in maternal performance increases with early-life fecundity in red deer.
 768 *Ecology Letters* 9, 1342-1350.

- 769 Oram, F., 2018. Abundance, behavioural and feeding ecology of wild orangutans (*Pongo*
 770 *pygmaeus morio*) in the fragmented forests of the Kinabatangan floodplain. Ph.D.
 771 Dissertation, Universiti Malaysia Sabah.
- 772 Palombit, R.A., 2012. Infanticide: male strategies and female counterstrategies. In: Mitani, J.C.,
 773 Call, J., Kappeler, P.M., Palombit, R.A., Silk, J.B. (Eds.), *The Evolution of Primate Societies*.
 774 Chicago University Press, Chicago, pp. 432-468.
- 775 Phillipps, Q., Phillipps, K., 2016. Phillipps' Field Guide to the Mammals of Borneo and Their
 776 Ecology: Sabah, Sarawak, Brunei, and Kalimantan. Princeton Field Guides, Princeton NJ.
- 777 Pontzer, H., Kamilar, J.M., 2009. Great ranging associated with greater reproductive investment
 778 in mammals. *Proceedings of the National Academy of Sciences USA* 106, 192-196.
- 779 Pontzer, H., Wrangham, R.W., 2004. Climbing and the daily energy cost of locomotion in wild
 780 chimpanzees: implications for hominoid locomotor evolution. *Journal of Human*
 781 *Evolution* 46, 315-333.
- 782 Pontzer, H., Brown, M.H., Raichlen, D.A., Dunsworth, H., Hare, B., Walker, K., Luke, A., Dugas,
 783 L.R., Durazo-Arvizu, R., Schoeller, D., Plange-Rhule, J., Bovet, P., Forrester, T.E., Lambert,
 784 E.V., Emery Thompson, M., Shumaker, R.W., Ross, S.R., 2016. Metabolic acceleration and
 785 the evolution of human brain size and life history. *Nature* 533, 390-392.
- 786 R-Core Team, 2016. R: A language and environment for statistical computing, 3.3.0. R
 787 Foundation for Statistical Computing, Vienna.
- 788 Reichard, U.H., Barelli, C., 2008. Life history and reproductive strategies of Khao Yai *Hylobates*
 789 *lar*: Implications for social evolution in apes. *International Journal of Primatology* 29,
 790 823-844

- 791 Rijksen, H.D., 1978. A Fieldstudy on Sumatran Orang Utans (*Pongo pygmaeus abelii* Lesson
792 1827). H. Veenman & Zonen, B.V., Wageningen.
- 793 Robson, S.L., van Schaik, C.P., Hawkes, K., 2006. The derived features of human life history. In:
794 Hawkes, K., Paine, R.R. (Eds.), The Evolution of Human Life History. School of American
795 Research Press, Santa Fe. pp. 17-44.
- 796 Schuppli, C., Forss, S.I.F., Meulman, E.J.M., Zweifel, N., Lee, K.C., Rukmana, E., Vogel, E.R., van
797 Noordwijk, M.A., van Schaik, C.P., 2016. Development of foraging skills in two orangutan
798 populations: needing to learn or needing to grow? *Frontiers in Zoology* 13, 43.
- 799 Shumaker, R.W., Wich, S.A., Perkins, L., 2008. Reproductive life history traits of female
800 orangtuan (*Pongo* spp.). In: Atsalis, S., Margulis, S.W., Hof, P.R. (Eds.), *Primate*
801 *Reproductive Aging*. Karger, Basel, pp. 147-161.
- 802 Sodaro, C., Frank, E., Nacey, A., Czekala, N., 2007. Orangutan development, reproduction and
803 birth management. In: Sodaro, C. (Ed.), *Orangutan Species Survival Plan Husbandry*
804 *Manual*. Chicago Zoological Society, Chicago, pp. 76-109.
- 805 Stearns, S.C., 2000. Life history evolution: successes, limitations, and prospects.
806 *Naturwissenschaften* 87, 476-486.
- 807 Terborgh, J., van Schaik, C.P., 1987. Convergence vs. non-convergence in primate communities.
808 In: Gee, J.H.R., Giller, P.S. (Eds.), *Organization of Communities: Past and Present*.
809 Blackwell Scientific Publications, Oxford, pp. 205-226.
- 810 Therneau, T.M., 2015a. A package for survival analysis in S. version 2.38. [https://CRAN.R-](https://CRAN.R-project.org/package=survival)
811 [project.org/package=survival](https://CRAN.R-project.org/package=survival)
- 812 Therneau, T.M., 2015b. coxme: Mixed Effects Cox Models. version 2.2-5. [https://CRAN.R-](https://CRAN.R-project.org/package=coxme)
813 [project.org/package=coxme](https://CRAN.R-project.org/package=coxme)

- 814 Thorpe, S.K.S., Crompton, R.H., 2006. Orangutan positional behavior and the nature of arboreal
815 locomotion in Hominoidea. *American Journal of Physical Anthropology* 131, 384-401.
- 816 van Noordwijk, M.A., van Schaik, C.P., 1999. The effects of dominance rank and group size on
817 female lifetime reproductive success in wild long-tailed macaques, *Macaca fascicularis*.
818 *Primates* 40, 105-130.
- 819 van Noordwijk, M.A., van Schaik, C.P., 2005. Development of ecological competence in
820 Sumatran orangutans. *American Journal of Physical Anthropology* 127, 79-94.
- 821 van Noordwijk, M.A., Arora, N., Willems, E.P., Dunkel, L.P., Amda, R.N., Mardianah, N.,
822 Ackermann, C., Krützen, M., van Schaik, C.P., 2012. Female philopatry and its social
823 benefits among Bornean orangutans. *Behavioral Ecology and Sociobiology* 66, 823-834.
- 824 van Noordwijk, M.A., Willems, E.P., Utami Atmoko, S.S., Kuzawa, C.W., van Schaik, C.P., 2013.
825 Multi-year lactation and its consequences in Bornean orangutans (*Pongo pygmaeus*
826 *wurmbii*). *Behavioral Ecology and Sociobiology* 67, 805-814.
- 827 van Schaik, C.P., 1999. The socioecology of fission-fusion sociality in orangutans. *Primates* 40,
828 73-90.
- 829 van Schaik, C.P., 2016. *The Primate Origins of Human Nature*. John Wiley & Sons, Hoboken, N.J.
- 830 van Schaik, C.P., Deaner, R.O., 2003. Life history and cognitive evolution in primates. In: de
831 Waal, F.B.M., Tyack, P.L. (Eds.), *Animal Social Complexity*. Harvard University Press,
832 Cambridge, MA., pp. 5-25.
- 833 van Schaik, C.P., Isler, K., 2012. Life-history evolution in primates. In: Mitani, J.C., Call, J.,
834 Kappeler, P.M., Palombit, R.A., Silk, J.B. (Eds.), *The Evolution of Primate Societies*.
835 University of Chicago Press, Chicago and London, pp. 220-244.

- 836 van Schaik, C.P., Pfannes, K.R., 2005. Tropical climates and phenology: a primate perspective. In:
 837 Brockman, D.K., van Schaik, C.P. (Eds.), *Seasonality in Primates: Studies of Living and*
 838 *Extinct Human and Non-Human Primates*. Cambridge University Press, Cambridge, pp.
 839 23-54.
- 840 van Schaik, C.P., Marshall, A.J., Wich, S.A., 2009. Geographic variation in orangutan behavior
 841 and biology. In: Wich, S.A., Utami Atmoko, S.S., Mitra Setia, T., van Schaik, C.P. (Eds.),
 842 *Orangutans: Geographic Variation in Behavioral Ecology and Conservation*. Oxford
 843 University Press, Oxford, pp. 351-361.
- 844 Vogel, E. R., Knott, C.D., Crowley, B.E., Blakely, M.D., Larsen, M.D., Dominy, N.J., 2012. Bornean
 845 orangutans on the brink of protein bankruptcy. Biology Letters 8: 333-336.
- 846 Walker, K.K., Walker, C.S., Goodall, J., Pusey, A.E., 2018. Maturation is prolonged and variable in
 847 female chimpanzees. *Journal of Human Evolution* 114, 131-140.
- 848 Walker, R., Gurven, M., Hill, K., Migliano, A.B., Chagnon, N., De Souza, R., Djurovic, G., Hames,
 849 R., Hurtado, M., Kaplan, H., Kramer, K., Oliver, W.J., Vallengia, C., Yamauchi, T., 2006.
 850 Growth rates and life histories in twenty-two small-scale societies. *American Journal of*
 851 *Human Biology* 18, 295-311.
- 852 Watts, D.P., 2012. The apes: Taxonomy, biogeography, life histories, and behavioral ecology. In:
 853 Mitani, J.C., Call, J., Kappeler, P.M., Palombit, R.A., Silk, J.B. (Eds.), *The Evolution of*
 854 *Primate Societies*. University of Chicago Press, Chicago and London, pp. 113-147.
- 855 Wich, S.A., Sterck, E.H.M., Utami, S.S., 1999. Are orang-utan females as solitary as chimpanzee
 856 females? *Folia Primatologica* 70, 23-28.

- 857 Wich, S.A., Utami-Atmoko, S.S., Mitra Setia, T., Rijksen, H.D., Schurmann, C., van Hooff,
858 J.A.R.A.M., van Schaik, C.P., 2004. Life history of wild Sumatran orangutans (*Pongo*
859 *abelii*). *Journal of Human Evolution* 47, 385-398.
- 860 Wich, S.A., de Vries, H., Ancrenaz, M., Perkins, L., Shumaker, R.W., Suzuki, A., van Schaik, C.P.,
861 2009a. Orangutan life history variation. In: Wich, S.A., Utami Atmoko, S.S., Mitra Setia,
862 T., van Schaik, C.P. (Eds.), *Orangutans: Geographic Variation in Behavioral Ecology and*
863 *Conservation*. Oxford University Press, Oxford, pp. 65-75.
- 864 Wich, S.A., Shumaker, R.W., Perkins, L., de Vries, H., 2009b. Captive and wild orangutan (*Pongo*
865 *sp.*) survivorship: A comparison and the influence of management. *American Journal of*
866 *Primatology* 71, 680-686.
- 867 Wilcox, C.H., Supiansyah, Abdul Azis, K., Zainuddin J., Rahman, Cheyne, S.M., 2016. Predator
868 mobbing and interspecies cooperation: an interaction between gibbons, langurs and a
869 clouded leopard. *Asian Primates Journal* 6, 20-26.
- 870 Wood, B.M., Watts, D.P., Mitani, J.C., Langergraber, K.E., 2017. Favorable ecological
871 circumstances promote life expectancy in chimpanzees similar to that of human hunter-
872 gatherers. *Journal of Human Evolution* 105, 41-56.

873 Table 1.

874 Data collection period per field site.

875

876 Table 2.

877 Classification criteria for infant birth dates and maternal stage.

878

879 Table 3

880 Interbirth interval between live births: the Cox proportional hazards models shows no significant

881 difference between subspecies (quadratic trend sequence *abelii-wurmbii-morio*) despite the longer 'tail'

882 for the Sumatran orangutan (*P. abelii*). Similarly, there is no significant effect of infant sex. The (NS)

883 quadratic trend for maternal stage indicates that multiparous females tend to have shorter interbirth

884 intervals.

885

886 Table 4

887 Cox proportional hazards model of offspring survival. The subspecies quadratic trend is based on

888 sequence *abelii-wurmbii-morio*, suggesting somewhat higher survival for *wurmbii* than either *abelii* or

889 *morio*. Similarly, offspring of multiparous females have higher survival than either young or old mothers.

890 No significant difference between males and female infants in survivorship.

Figure legends

Figure 1

Location of study sites: *Pongo abelii* in 1. Suaq Balimbing and 2. Ketambe; *P. pygmaeus wurmbii* in 3. Gunung Palung, 4. Sebangau and 5. Tuanan; and *P.p. morio* in 6. Danum Valley and 7. Lower Kinabatangan.

Figure 2

Kaplan –Meier curves for age of first reproduction (AFR), $n=45$ females (with birth date known within 6 months), including four females who gave birth for the first time and 41 who had not given birth by the end of the study (right-censored values). The right-hand graph shows there are too few completed AFRs to detect an effect of maternal age on her daughter's AFR.

Figure 3

Kaplan-Meier curves for interbirth intervals $n= 99$, with 49 completed intervals.

a) Comparison between islands; b) Comparison between subspecies; c) Comparison of interbirth intervals after surviving male or female offspring, d) Comparison between females of different maternal stage.

Note: Excluded from calculations were two old females without new offspring for >14 years since their last known one.

Figure 4

Interval between death of an unweaned offspring and the next birth, in relation to the age at which the previous infant died. The linear trend for the Bornean data is indicated by a dashed line.

Figure 5

Kaplan-Meier curves for survival of immatures: a) Comparison between islands; b) Comparison between subspecies; c) Comparison of male or female offspring, d) Comparison of offspring of females of different maternal stage. 108 individuals from seven sites, belonging to three subspecies

921
922 Figure 6
923 Comparative survival until 18 years among female great apes and modern human populations.
924 The box plot with median (bar) and mean (x) values at 15 years is based on the average
925 survivorship in 13 small-scale societies taken from Walker et al. (2006). The orangutan curve is
926 surrounded by its confidence limits.

927 Sources: orangutan (wild) - this study; orangutan (captive) – Wich et al., 2009a; Swiss
928 population-Schweizerische Bundesamt für Statistik, 2017; Hadza – Blurton-Jones, 2016; Ngogo
929 chimpanzees - Wood et al., 2017; Kanyawara chimpanzees – Muller and Wrangham, 2014;
930 gorilla – Bronikowski et al., 2016a, b.

931 Table 1

932 Data collection period per field site.

933

Island	Site	Start data	Last data	Max. span of study in months	Coordinating researcher
Sumatra: <i>Pongo abelii</i>					
<i>P. abelii</i>	Ketambe - reintro, Aceh tenggara*/**	1972	June 2015	516	Sri Suci Utami Atmoko
	Ketambe - wild, Aceh tenggara*	1972	June 2015	516	Sri Suci Utami Atmoko
	Suaq Balimbing, Aceh selatan *	Jan. 1994 Jul. 2007	Sept. 1999 Feb. 2017	277	Carel van Schaik; Caroline Schuppli
Borneo: <i>Pongo pygmaeus</i>					
<i>P. p. wurmbii</i>	Gunung Palung, West Kalimantan *	1994 2008	2003 June 2016	252	Cheryl Knott Cheryl Knott
	Sabangau, Central Kalimantan	2003	May 2016	156	Helen Morrogh- Bernard
	Tuanan, Central Kalimantan	June 2003	Feb. 2017	164	Maria van Noordwijk
	KOCP-Sukau	1998	Dec. 2016	221	Felicity Oram
<i>P. p. morio</i>	Danum Valley, Sabah	2004	Dec. 2016	149	Noko Kuze

* monitoring interrupted for multiple years due to external circumstances

** data only included in SOM Table S2

934

935 Table 2

936 Classification criteria for infant birth dates and maternal stage.

937 **Infant birth date**

Class	Accuracy	Criteria to assess birth date
known	within 1 month	mother known and observed before and after birth of offspring
estim.	within 3 months	mother known and observed before and after birth of offspring: "estimate"
rough estim.	within 6 months	mother known and observed before and after birth of offspring, or infant extremely young at first encounter of the mother: "rough estimate"
very rough estim.	within 1 year	estimated age for dependent offspring born before first encounter of mother, but estimated <3 y old: "very rough estimate"
unknown	age & year guessed	estimated age for dependent offspring born before first encounter of mother, but estimated at least 3 y old

Maternal stage

Class	Criteria to assess maternal stage
young	first offspring and < 21 years - thus if first infant lost pre-weaning and second one born when <21 years, still counted as young
multiparous	≥ 21 years and 2 nd - 4 th birth
old	birth 5 or later and estimated >35 year

938

Table 3

Interbirth interval between live births: the Cox proportional hazards model shows no significant difference between subspecies (quadratic trend sequence *abelii-wurmbii-morio*) despite the longer ‘tail’ for the Sumatran (*P. abelii*). Similarly, there is no significant effect of infant sex. Multiparous females tend to have shorter interbirth interval (NS).

Cox proportional hazard mixed-effects model Interbirth interval

	B	se	Hazard ratio (95% CI)	z	p
Subspecies					
Linear trend	-0.320	0.76	0.726 (0.165 – 3.192)	-0.42	0.670
Quadratic trend	-0.865	0.60	0.421 (0.131 – 1.354)	-1.45	0.150
Infant sex					
Female	-	-	-	-	-
Male	0.140	0.52	1.151 (0.417 – 3.176)	0.27	0.790
Maternal stage					
Linear trend	-0.858	0.73	0.424 (0.101 – 1.771)	-1.18	0.240
Quadratic trend	-0.890	0.49	0.411 (0.158 – 1.068)	-1.82	0.068

57 females from 7 sites, belonging to 3 subspecies

$n_{\text{births}} = 99$, $n_{\text{life-life IBI}} = 49$, $\chi^2_{\text{LRT}} = 5.32$, $p = 0.379$

Table 4

Cox proportional hazards model of offspring survival. The subspecies quadratic trend is based on sequence *abelii-wurmbii-morio*, suggesting somewhat higher survival for *wurmbii* than either *abelii* or *morio*. Similarly, offspring of multiparous females have higher survival than either young or old mothers. No significant difference between males and female infants in survivorship.

Cox proportional hazard mixed-effects model offspring survival

	B	se	Hazard ratio (95% CI)	z	p
Subspecies					
Linear trend	1.101	0.61	3.006 (0.910 – 9.930)	1.81	0.071
Quadratic trend	1.198	0.60	3.314 (1.018 – 10.785)	1.99	0.047
Infant sex					
Female	-	-	-	-	-
Male	1.384	0.77	3.990 (0.853 – 18.648)	1.76	0.079
Maternal stage					
Linear trend	0.031	0.53	1.031 (0.364 – 2.920)	0.06	0.950
Quadratic trend	1.796	0.70	6.027 (1.520 – 23.905)	2.56	0.011

108 individuals from 7 sites, belonging to 3 subspecies

$n_{\text{individuals}} = 108$, $n_{\text{deaths}} = 10$, $\chi^2_{\text{LRT}} = 14.21$, $p = 0.014$

960 Table 5

961 Orangutan female life-table (0-18 years)

Age (years)	Enter	Censored	Dead	At Risk	q_x	p_x	l_x
0-1	52	1	1	51	0.0196	0.9804	1.0000
1-2	50	3	1	47	0.0213	0.9787	0.9804
2-3	46	6	1	40	0.0250	0.9750	0.9595
3-4	39	5	1	34	0.0294	0.9706	0.9355
4-5	33	3	0	30	0.0000	1.0000	0.9080
5-6	30	6	0	24	0.0000	1.0000	0.9080
6-7	24	0	0	24	0.0000	1.0000	0.9080
7-8	24	4	0	20	0.0000	1.0000	0.9080
8-9	20	1	0	19	0.0000	1.0000	0.9080
9-10	19	6	0	13	0.0000	1.0000	0.9080
10-11	13	4	0	9	0.0000	1.0000	0.9080
11-12	9	3	0	6	0.0000	1.0000	0.9080
12-13	6	1	0	5	0.0000	1.0000	0.9080
13-14	5	2	0	3	0.0000	1.0000	0.9080
14-15	3	0	0	3	0.0000	1.0000	0.9080
15-16	3	1	0	2	0.0000	1.0000	0.9080
16-17	2	1	0	1	0.0000	1.0000	0.9080
17-18	1	1	0	0	0.0000	1.0000	0.9080

962

963

964

965

966

967 Supplementary Online Material

968 Tables

969

970 The slow ape: high infant survival and long interbirth intervals in wild orangutans

971

972 Maria A. van Noordwijk¹, S. Suci Utami Atmoko², Cheryl D. Knott³, Noko Kuze⁴, Helen C.

973 Morrogh-Bernard⁵, Felicity Oram⁶, Caroline Schuppli¹, Carel P. van Schaik¹, Erik P. Willems¹

974 Table S1

975 Cox model for female age of first reproduction, using least restrictive criteria of inclusion (very
976 rough estimates). The subspecies quadratic trend is based on sequence *abelii-wurmbii-morio*.

977

	B	se	Hazard ratio (95% CI)	z	p
Subspecies					
Linear trend	-0.509	0.86	0.601 (0.112 - 3.232)	-0.59	0.55
Quadratic trend	-0.165	0.62	0.848 (0.250 - 2.879)	-0.27	0.79

58 females from 7 sites, belonging to 3 subspecies

$n_{\text{individuals}} = 58$, $n_{\text{events}} = 13$, $\chi^2_{\text{LRT}} = 0.36$, $p = 0.834$

978

979 Table S2

980 Cause of death of immatures. (Unless a reference is given the details of these cases are

981 unpublished – please refer to contributing author per site, see Table 1).

982

Site	Mother	Infant	Age death in months	Stage	Cause of death
Danum V	Yanti	Jun	c 72	post-Wean	prob. clouded leopard ¹
Danum V	Yanti	Toy	53	pre-Wean	?
Gn. Pal.	Walimah	Wanda	4	pre-Wean	mother wounded: maybe infanticide ²
Sabangau	Indah	Indah-inf 3	3	pre-Wean	in rain storm
Low Kin	Jenny	Dhani	0,25	pre-Wean	newborn-bad health
Low Kin	Mawar	Lara	36	pre-Wean	?
Tuanan	Kondor	Kahiyu	26	pre-Wean	?
Tuanan	Kondor	Kilmino	18	pre-Wean	?
Tuanan	Sumi	Susi	48	pre-Wean	after mother died of wounds - prob. by clouded leopard ³
Suaq	Friska	Henki	18	pre-Wean	?
Suaq	Tevi	Tedi	36	pre-Wean	?
Suaq	Butet	B	12	pre-Wean	mother bad health
<i>Additional cases not included in this study</i>					
DV	Lina	Lina-inf	0	pre-Wean	?
Tuanan	Pinky	Pina	15	pre-Wean	mother survived, but wounded - prob. by clouded leopard
Suaq	Ani	Aneka	<132		?
<i>Released orangutans and descendants in Ketambe</i>					
Ket-R	Binjei	Harto	84	post-W	?
Ket-R	Ans	Ari	6	pre-Wean	after mother died
Ket-R	Binjei	Dona	14	pre-Wean	after mother died
Ket-R	Getti	Gendhuk	9	pre-Wean	after mother died

983 ¹Kanamori et al., 2012; ²C.D. Knott et al., unpublished

984 Table S3

985 Selected female survival values from human population in the 20th century in Switzerland and
 986 USA.

987

Country	Year	L _x at 15 years	Reference
Switzerland	1900	80,6	Bundesamt für Statistik, Kohortensterbetafel für die Schweiz (1900-2030) nach Geburtsjahrgang, Geschlecht und Alter. Accessed 26 Sep 2017 https://www.bfs.admin.ch/bfs/de/home/statistiken/bevoelkerung
Switzerland	1925	91,9	See above
Switzerland	1950	96,4	See above
Switzerland	2000	99,4	See above
USA	1900	78,3	F.C. Bell and M.L. Miller Life Tables for the United States of America Social Security Area 1900-2100 Actuarial Study No. 116. accessed through Human Life-Table Database, http://www.lifetable.de
USA	1925	90,5	See above
USA	1950	96,5	See above
USA	2000	99,1	Arias, E., United States Life Tables, 2000, National Vital Statistics Reports, Volume 51-3, 2002. accessed through Human Life-Table Database, http://www.lifetable.de

988

989 Table S4

990 Site-specific interbirth intervals.

991

Site	Closed intervals with previous infant alive				Closed intervals with previous infant alive and evidence of miscarriage			
	Mean IBI (years)	Median IBI (years)	Range (months)	<i>n</i>	Mean IBI (years)	Median IBI (years)	Range (months)	<i>n</i>
BORNEO								
Danum Valley	8.5	8.5	84-102	2	7.0	7.0	84	1
Gunung Palung	7.9	7.6	82-124	11	7.6	7.6	82-99	10
Kinabatangan	7.4	7.5	78-96	4	7.4	7.5	78-96	4
Sebangau	7.2	7.25	66-107	5	6.8	7.0	66-94	4
Tuanan	7.0	7.0	71-100	13	7.0	7.0	71-100	12
SUMATRA								
Ketambe	8.5	7.75	67-168	16	7.7	7.25	67-168	14
Suaq Balimbing	7.5	7.2	75-112	8	7.5	7.2	75-112	8

992

993 Table S5

994 Site-specific infant mortality and age at first birth (AFR). * AFR with at least roughly estimated
 995 dates of own birth and first infant's birth (both < +/- 6 months), all other AFRs are very rough
 996 estimates (at least one of the values \pm 6-12 months).

997

Site	Mean % infant mortality	<i>n</i> (births)	Mean AFR (years)	Range AFR (years)	<i>n</i> (females)
BORNEO					
Danum Valley	13	16			0
Gunung Palung	4	28	15.1	14.4-16.4*	3
Kinabatangan	12	19	15.5		1
Sebangau	5	19	13.3	12.7-13.8	2
Tuanan	9	35	14.1	13.3-15.0	2
SUMATRA					
Ketambe	0	32	13.7	13.3*-14.1*	2
Suaq Balimbing	15	27	14.8	13.7-15.7*	3

998

999

1000

Figure 1



Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

